

# The gaussian image of mean curvature one surfaces in $\mathbb{H}^3$ of finite total curvature.

Pascal Collin      Laurent Hauswirth      Harold Rosenberg

ABSTRACT.<sup>1</sup>- The hyperbolic Gauss map  $G$  of a complete constant mean curvature one surface  $M$  in hyperbolic 3-space, is a holomorphic map from  $M$  to the Riemann sphere. When  $M$  has finite total curvature, we prove  $G$  can miss at most three points unless  $G$  is constant. We also prove that if  $M$  is a properly embedded mean curvature one surface of finite topology, then  $G$  is surjective unless  $M$  is a horosphere or catenoid cousin.

We consider complete surfaces  $M$  in hyperbolic 3-space  $\mathbb{H}^3$  with mean curvature one and of finite total curvature. For a point  $q \in M$ , the Gauss map  $G$  sends  $q$  to the point at infinity obtained as the positive limit of the geodesic of  $\mathbb{H}^3$  starting at  $q$  and having  $\vec{H}(q)$  (the mean curvature vector of  $M$  at  $q$ ) as its tangent at  $q$ . Bryant has shown that  $G$  is meromorphic on  $M$  and  $M$  admits a parametrization by meromorphic data analogous to the Weierstrass representation of minimal surfaces in Euclidean 3-space  $\mathbb{R}^3$  [?], [?].

Yu [?] has shown that  $G$  can omit at most 4 points of the sphere at infinity  $S_\infty$ , unless  $M$  is a horosphere and  $G$  is constant. For complete minimal surfaces in  $\mathbb{R}^3$  of finite total curvature, Osserman had shown that the Gauss map omits at most 3 points of the sphere, unless  $M$  is a plane. In this paper we establish a result of this type in  $\mathbb{H}^3$ .

The conformal type of a complete surface of mean curvature one with finite total curvature in  $\mathbb{H}^3$  is finite, i.e.,  $M$  is conformally a compact Riemann surface  $\bar{M}$  with a finite number of points removed (called the punctures), but  $G$  does not necessarily extend meromorphically to the punctures.  $M$  is called regular when  $G$  does extend meromorphically to the punctures.

---

<sup>1</sup>MSC : 53A10 (53A35)

Our first result is then:

**Theorem 1** *Let  $M$  be a complete surface immersed in  $\mathbb{H}^3$  with mean curvature one and of finite total curvature. Then  $G$  can omit at most 3 points unless  $G$  is constant and  $M$  is a horosphere.*

*Proof.* If  $G$  is not regular, then  $G$  has an essential singularity at a puncture  $p_0$ . By Picard's theorem,  $G$  can omit at most two values in a neighborhood of this puncture. Thus in the following we can assume that  $G$  is meromorphic on  $\overline{M}$ , i.e.,  $M$  is regular.

Let  $(g, \omega)$  be local Weierstrass data of the minimal cousin of  $M$  in  $\mathbb{R}^3$  (cf. [?], [?] for the details). The induced metric on  $M$  is given by  $ds = (1 + |g|^2)|\omega|$ , and the holomorphic quadratic differential

$$Q = \omega dg$$

is globally defined on  $M$  and meromorphic at each puncture of  $M$ , with a pole at each puncture which is at worst of order 2. Since  $dG$  is meromorphic on  $\overline{M}$  (the conformal compactification of  $M$ ), the 1-form  $\omega^\# = Q/dG$  is meromorphic on  $\overline{M}$ ; in a local conformal coordinate,  $\omega^\# = (g'(z)/G'(z))\omega(z)$ .

The Schwarzian quadratic differentials of  $g$ ,  $G$  and  $Q$  are related on  $\overline{M}$  ([?], [?]):

$$S(g) - S(G) = 2Q, \tag{1}$$

where  $S(g)(z) = \left( \left( \frac{g''}{g'} \right)' - \frac{1}{2} \left( \frac{g''}{g'} \right)^2 \right) dz^2$ . Writing  $g(z) = a_0 + z^k(a_1 + a_2z + \dots)$ , a calculation shows that  $S(g)$  has at worst a pole of order 2 at  $z$  and the coefficient of  $dz^2/z^2$  is  $(1 - k^2)/2$ .

Since  $Q$  is holomorphic on  $M$ , it follows from (??) that the branch points and non-simple poles of  $g$  and  $G$  on  $M$  coincide with each other and each of them has the same multiplicity (the branching order of  $g$  at  $z$  is defined to be  $k - 1 = b_g(z)$ ). In particular,  $\omega^\#$  has no poles on  $M$ .

We next observe that the zeros of  $\omega^\#$  on  $M$  are the poles of  $G$  on  $M$ , and a pole of  $G$  of order  $k$  is a zero of  $\omega^\#$  of order  $2k$ . First, suppose that  $z \in M$  is a pole of  $G$  of order  $k$ . Then  $k \geq 1$  and  $z$  may, or may not, be a pole of  $g$ . If it is a pole of  $g$ , then  $z$  is a pole of  $g$  of order  $k$  (by the Schwarzian derivative relation) and then is a zero of  $\omega$  of order  $2k$ . Hence the order of a zero of  $\omega^\#$  is of twice the order of the pole of  $G$ . If  $z$  is not a pole of  $g$ , then it is not a zero of  $\omega$  but a zero of  $g'$  of order

$k - 1$  and a pole of  $G'$  of order  $k + 1$ . Consequently  $\omega^\#$  also has a zero whose order is twice the order of the pole of  $G$ . An analogous computation, in the case that  $G$  has no poles, implies that  $\omega^\#$  is holomorphic and not zero.

Let  $p_1, \dots, p_r$  be the punctures, so  $\overline{M} = M \cup \{p_1, \dots, p_r\}$ . After an isometry of  $\mathbb{H}^3$ , we can suppose that  $G$  has only simple poles on  $M$  and has no zeros or poles at the punctures. The metric

$$ds^\# = (1 + |G|^2)|\omega^\#|$$

is complete on  $\overline{M}$ , so  $\omega^\#$  has a pole at each puncture [?]. The order of the pole of  $\omega^\#$  at  $p_j$  is given by

$$P_{p_j}(\omega^\#) = \lambda_Q(p_j) + b_G(p_j),$$

where  $Q(z) = (\gamma/(z - p_j)^{\lambda_Q(p_j)} + \dots)dz^2$  is the Laurent expansion of  $Q$  at  $p_j$ . Then the total order of the poles of  $\omega^\#$  is

$$P(\omega^\#) = \sum_{j=1}^r \lambda_Q(p_j) + \sum_{j=1}^r b_G(p_j). \quad (2)$$

By Riemann's relation for  $\omega^\#$  on  $\overline{M}$ , we have

$$P(\omega^\#) - 2N = 2 - 2s, \quad (3)$$

where  $N$  is the degree of  $G$  (so  $2N$  is the order of zeros of  $\omega^\#$ , since  $G$  has  $N$  simple poles on  $M$ ) and  $s$  is the genus of  $M$ .

Let  $q_1, \dots, q_k$  be the points of  $S_\infty$  omitted by  $G$ , so that  $G^{-1}\{q_1, \dots, q_k\} \subset \{p_1, \dots, p_r\}$  (we write  $G$  also for the meromorphic extension of  $G$  to  $\overline{M}$ ). Then we have

$$kN \leq \sum_{j=1}^r (1 + b_G(p_j)) \leq r + b, \quad (4)$$

where  $b$  is the total branching order of  $G$ . Here  $1 + b_G(p_j)$  is the total number of times that  $G$  takes its value at  $p_j$ , counted with multiplicity.

Riemann's relation applied to the 1-form  $dG$  on  $\overline{M}$  yields:

$$2N - b = 2 - 2s. \quad (5)$$

Now by Lemma 3 of [?], we have at each puncture  $p_j$ :

$$\lambda_Q(p_j) + b_G(p_j) \geq 2.$$

Then equation (??) gives:

$$P(\omega^\#) \geq 2r. \tag{6}$$

This last inequality together with the equations (??) and (??) yields:

$$P(\omega^\#) = 4N - b \geq 2r.$$

Then the equation (??) implies:

$$4N - kN \geq r \geq 1, \tag{7}$$

and  $k$  is at most 3. □

**Theorem 2** *Let  $M$  be a properly embedded surface in  $\mathbb{H}^3$  with mean curvature one and of finite topology. If  $M$  is not a horosphere nor a catenoid cousin, then the Gauss map  $G$  of  $M$  is surjective.*

*Proof.* We know that  $M$  has finite total curvature and each end of  $M$  is regular [?]; also each end is asymptotic to an end of a horosphere or an end of a catenoid cousin. We also proved in [?] that the asymptotic boundary of an end is precisely the limiting value of  $G$  at the puncture. We can suppose  $M$  has at least two ends, since if  $M$  had only one end, the asymptotic boundary of  $M$  would be one point and  $M$  would be a horosphere [?].

We claim that each end of  $M$  is asymptotic to a catenoid cousin end. Suppose this were not true. Let  $E$  be an end of  $M$  asymptotic to a horosphere end. We work in the upper half-space model of  $\mathbb{H}^3$ ,  $\{x_3 > 0\}$ , and assume  $E$  is asymptotic to a horosphere  $x_3 = c > 0$ . In particular, the mean curvature vector of  $E$  points up outside of some compact set of  $E$ . There are no ends of  $M$  above  $E$ . Indeed, their mean curvature vector would also point up (each such end is asymptotic to a horizontal horosphere or a catenoid cousin end whose limiting normal points vertically up) and  $M$  separates  $\mathbb{H}^3$  into two connected components, so no such end is above  $E$ .

Then for  $\varepsilon > 0$ , the part  $A$  of  $M$  above  $c + \varepsilon$  is compact. At the highest point of  $A$  (if  $A$  were not empty) the mean curvature vector of  $M$  points down. But this highest point can be joined by an arc in  $\mathbb{H}^3 \setminus M$ , to a point of  $E$  where the mean curvature vector points up. Thus  $M$  is completely below  $x_3 = c$ .

Let  $\varepsilon > 0$ , and let  $C$  be a small circle in the plane  $x_3 = c - \varepsilon$  so that  $C$  is above  $M$ . Just as in the proof of the half-space theorem for properly immersed minimal surfaces in  $\mathbb{R}^3$  [?], one can take a family of catenoid cousin ends  $C(\lambda)$ ,  $\partial C(1) = C$

with  $C(1)$  above  $M$ , and  $C(\lambda)$  converges to the plane  $x_3 = c - \varepsilon$  as  $\lambda \rightarrow 0$ . Then some  $C(\lambda)$  would touch  $M$  at a point  $q \in M$ , and the maximum principle would yield  $M$  equals this catenoid cousin. Thus each end of  $M$  is asymptotic to a catenoid cousin.

Next, observe that  $G$  is injective on the set of punctures; two distinct ends can not be asymptotic to the same point at infinity. This follows from the fact that each end is asymptotic to a catenoid cousin end and we know the direction of the mean curvature vector along the end. When  $M$  is embedded,  $M$  separates  $\mathbb{H}^3$  and the mean curvature vector points into one of the components of the complement. Thus two ends can not be asymptotic to the same point at infinity.

Now, suppose that  $G$  is not surjective and omits a point  $q$ . Then there is exactly one catenoid cousin type end  $E$  of  $M$  asymptotic to  $q$ . Let  $p \in \bar{M}$  be the puncture of  $E$  such that  $G(p) = q$ . We know  $G$  has local degree one at  $p$ . There is no other point  $p' \in \bar{M}$  sent to  $q$  by  $G$ . For  $p'$  can not be a puncture of  $M$ , since  $G$  is injective on the punctures, and  $p'$  can not be a point of  $M$  because  $q$  is a value omitted. Hence the degree  $N$  of  $G$  on  $\bar{M}$  is one.

We use the same notation as in Theorem 1. At each puncture  $p_j$  of  $M$ ,  $\omega^\#$  has a pole exactly of order 2. So, by equation (??), we have

$$2r - 2 = 2 - 2s \text{ and } r + s = 2.$$

Then  $M$  is the catenoid cousin ( $r = 2$ ) and Theorem 2 is proved. □

## References

- [1] R.Bryant, *Surfaces of mean curvature one in hyperbolic space*, Astérisque No.154-155 (1987), Soc. Math France, 321-347, 1988.
- [2] P.Collin, L.Hauswirth and H.Rosenberg, *The geometry of finite topology Bryant surfaces*, to appear in Ann. of Math.
- [3] D.Hoffman and W.Meeks, *The strong half-space theorem for minimal surfaces*, Invent. Math. **101** (1990), 373-377.
- [4] M.Umehara and K.Yamada, *Complete surfaces of constant mean curvature-1 in the hyperbolic 3-space*, Ann. of Math. **137** (1993), 611-638.

- [5] M.Umehara and K.Yamada, *A duality on CMC-1 surface in the hyperbolic 3-space and a hyperbolic analogue of the Osserman Inequality*, Tsukuba J. Math. **21** (1997), 229-237.
- [6] Zu-Huan Yu, *The value distribution of the hyperbolic Gauss map*, Proc. Amer. Math. Soc. **125** (1997), 2997-3001.

PASCAL COLLIN

UNIVERSITÉ PAUL SABATIER  
118, ROUTE DE NARBONNE  
31062 TOULOUSE - FRANCE  
*collin@picard.ups-tlse.fr*

LAURENT HAUSWIRTH

UNIVERSITÉ MARNE-LA-VALLÉE  
CITÉ DESCARTES  
5, BOULEVARD DESCARTES  
77454 CHAMPS-SUR-MARNE  
MARNE-LA-VALLÉE - FRANCE  
*hauswirth@math.univ-mlv.fr*

HAROLD ROSENBERG

UNIVERSITÉ DE PARIS 7  
2 PLACE JUSSIEU  
75005 PARIS - FRANCE  
*rosen@math.jussieu.fr*