

Pinching of the First Eigenvalue of the Laplacian and almost-Einstein Hypersurfaces of the Euclidean Space

Julien Roth

Institut Élie Cartan, UMR 7502
Nancy-Université, CNRS, INRIA
B.P. 239, 54506 Vandœuvre lès Nancy Cedex, France

roth@iecn.u-nancy.fr

Abstract

In this paper, we prove new pinching theorems for the first eigenvalue $\lambda_1(M)$ of the Laplacian on compact hypersurfaces of the Euclidean space. These pinching results are associated with the upper bound for $\lambda_1(M)$ in terms of higher order mean curvatures H_k . We show that under a suitable pinching condition, the hypersurface is diffeomorphic and almost isometric to a standard sphere. Moreover, as a corollary, we show that a hypersurface of the Euclidean space which is almost Einstein is diffeomorphic and almost isometric to a standard sphere.

Key words: Laplacian, eigenvalues, pinching, hypersurfaces, r -th mean curvatures, almost-Einstein

Mathematics Subject Classification: 53A07, 53C20, 53C21, 58C40.

1 Introduction and Preliminaries

Let (M^n, g) be a n -dimensional compact, connected, oriented manifold without boundary, isometrically immersed by ϕ into the $(n + 1)$ -dimensional Euclidean space, (\mathbb{R}^{n+1}, can) , *i.e.*, $\phi^*can = g$. If, in addition, (M^n, g) is Einstein, then a well-known result of Cartan and Thomas ([10]), also proved by Fialkow ([3]), says that M is a round sphere $\mathbb{S}^n(R)$ of corresponding radius.

A natural question is to ask what one could say if (M^n, g) is almost-Einstein, that is, $\|\text{Ric} - kg\|_\infty \leq \varepsilon$, for some positive constant k .

Recently, J.F. Grosjean gave a new proof of the Thomas-Cartan theorem using an upper bound of the first eigenvalue of the Laplacian. Indeed, Grosjean proved in [4] that if (M^n, g) has positive scalar curvature, then the first eigenvalue of the Laplacian satisfies

$$\lambda_1(M) \leq \frac{1}{n-1} \|\text{Scal}\|_\infty,$$

with equality only for geodesic spheres (here Scal denotes the scalar curvature).

If (M^n, g) is Einstein, *i.e.*, $\text{Ric} = (n-1)g$, we know by the Lichnerowicz theorem that $\lambda_1(M) \geq n$, and by the above upper bound

$$\lambda_1(M) \leq \frac{1}{n-1} \|\text{Scal}\|_\infty = n.$$

So $\lambda_1(M) = n$ and we are in the equality case of both inequalities, that is, $M = \mathbb{S}^n$.

This approach leads naturally to consider a pinching result on the first eigenvalue of the Laplacian, which allows to show that an almost Einstein hypersurface of \mathbb{R}^{n+1} is close to a sphere.

First, we can deduce from a theorem of Aubry ([1]), which is a pinching theorem corresponding to the Lichnerowicz inequality, that if ε is small enough, then M is homeomorphic to \mathbb{S}^n (see Theorem 3).

Nethertheless, Aubry's result does not yield to a sufficiently strong rigidity result. For this, we will study another pinching of the first eigenvalue of the Laplacian, which is associated with an extrinsic upper bound involving the scalar curvature. In fact, in this paper, we are interested in more general upper bounds in terms of higher order mean curvatures.

In [7], Reilly gives upper bounds for $\lambda_1(M)$, in terms of higher order mean curvatures H_k , which are defined as the symmetric polynomials in the principal curvatures. He shows that for all $k \in \{1, \dots, n\}$:

$$(1) \quad \lambda_1(M) \left(\int_M H_{k-1} \right)^2 \leq \frac{n}{V(M)} \int_M H_k^2,$$

with equality only for the standard spheres of \mathbb{R}^{n+1} .

By the Hölder inequality, we get similar inequalities with the L^{2p} -norms ($p \geq 1$) of H_k :

$$(2) \quad \lambda_1(M) \left(\int_M H_{k-1} \right)^2 \leq \frac{n}{V(M)^{1/p}} \|H_k\|_{2p}^2.$$

As for inequality (1), the equality case in (2) characterizes the standard spheres.

Then, a natural question is to know if there exists a pinching result as the following theorem proved by B. Colbois and J.F. Grosjean ([2])? *For $p \geq 2$ and any $\varepsilon > 0$, there exists a constant C_ε depending only on n and $\|H\|_\infty$ so that if the pinching condition*

$$\frac{n}{V(M)^{1/p}} \|H\|_{2p}^2 - C_\varepsilon < \lambda_1(M)$$

is true, then the Hausdorff distance between M and the sphere $S\left(0, \sqrt{\frac{n}{\lambda_1(M)}}\right)$ is at most ε .

We give a positive answer to this question, and, as we will see, the case $k = 2$, that is involving H_1 and H_2 , will solve the problem for almost-Einstein hypersurfaces.

Theorem 1. *Let (M^n, g) be a compact, connected, oriented Riemannian manifold without boundary isometrically immersed in \mathbb{R}^{n+1} and p_0 the center of mass of M . Assume that $V(M) = 1$ and let $k \in \{1, \dots, n\}$ such that $H_k > 0$. Then, for any $p \geq 2$ and for any $\varepsilon > 0$, there exists a constant C_ε depending only on ε , n , $\|H\|_\infty$ and $\|H_k\|_{2p}$ such that*

$$(P_{C_\varepsilon}) \quad \lambda_1(M) \left(\int_M H_{k-1} \right)^2 - \frac{n}{V(M)^{1/p}} \|H_k\|_{2p}^2 > -C_\varepsilon$$

is satisfied, then

$$i) \quad \phi(M) \subset B\left(x_0, \sqrt{\frac{n}{\lambda_1(M)}} + \varepsilon\right) \setminus B\left(x_0, \sqrt{\frac{n}{\lambda_1(M)}} - \varepsilon\right).$$

$$ii) \quad \forall x \in S\left(x_0, \sqrt{\frac{n}{\lambda_1(M)}}\right), \quad B(x, \varepsilon) \cap \phi(M) \neq \emptyset.$$

We recall that the Hausdorff distance between two compact subsets A and B of a metric space (E, d) is given by

$$d_H(A, B) = \inf \{ \eta > 0 \mid B \subset V_\eta(A) \text{ and } A \subset V_\eta(B) \},$$

where for any subset A , the set $V_\eta(A)$ is the tubular neighborhood of A defined by $V_\eta(A) = \{x \in E \mid d(x, A) < \eta\}$. So, *i*) and *ii*) of Theorem 1 imply that the Hausdorff distance between M and $S\left(x_0, \sqrt{\frac{n}{\lambda_1(M)}}\right)$ is at most ε .

Remark 1. We will see in the proof that $C_\varepsilon \longrightarrow 0$ when $\|H\|_\infty \longrightarrow \infty$ or $\varepsilon \longrightarrow 0$.

In this second theorem, if the pinching condition is strong enough, with a control on the L^∞ -norm of the second fundamental form B , we obtain that M is diffeomorphic and almost-isometric to a round sphere in the following sense

Theorem 2. Let (M^n, g) be a compact, connected, oriented Riemannian manifold without boundary isometrically immersed in \mathbb{R}^{n+1} and p_0 the center of mass of M . Assume that $V(M) = 1$ and let $k \in \{1, \dots, n\}$ such that $H_k > 0$. Then for any $p \geq 2$, there exists a constant K depending only on n , $\|B\|_\infty$ and $\|H_k\|_{2p}$ such that if the pinching condition

$$(P_K) \quad \lambda_1(M) \left(\int_M H_{k-1} \right)^2 - \frac{n}{V(M)^{1/p}} \|H_k\|_{2p}^2 > -K$$

is satisfied, then M is diffeomorphic to \mathbb{S}^n .

More precisely, there exists a diffeomorphism F from M into the sphere $\mathbb{S}^n \left(\sqrt{\frac{n}{\lambda_1(M)}} \right)$ of radius $\sqrt{\frac{n}{\lambda_1(M)}}$ which is a quasi-isometry. Namely, for any $\theta \in]0, 1[$, there exists a constant K_θ depending only on θ , n , $\|B\|_\infty$ and $\|H_k\|_{2p}$ so that the pinching condition with K_θ implies

$$| |dF_x(u)|^2 - 1 | \leq \theta,$$

for any unitary vector $u \in T_x M$.

Remark 2. We will see in the proof that the constants C_ε , K and K_θ of Theorems 1 and 2 do not depend on $\|H_k\|_{2p}$ if $p \geq \frac{n}{2k}$.

These results have a double interest. First, they improve the results in [2]. Second, the case $k = 2$ is especially interesting. Indeed, for hypersurfaces of the Euclidean space, the second mean curvature H_2 is, up to a multiplicative constant, the scalar curvature. Precisely, $H_2 = \frac{1}{n(n-1)\text{Scal}}$. Then we deduce from Theorems 1 and 2 two corollaries for almost-Einstein hypersurfaces.

Now, we give some preliminaries for the proof of these theorems. Throughout this paper, we consider a compact, connected, oriented Riemannian manifold isometrically immersed in $(\mathbb{R}^{n+1}, \text{can})$ by ϕ . Let ν be the outward normal unitary vector field. We denote respectively by ∇ and Δ the Riemannian connection and the Laplacian of M , and by $\bar{\nabla}$ the Riemannian connection

of \mathbb{R}^{n+1} . Finally, we denote by $\langle \cdot, \cdot \rangle$ the Euclidean scalar product of \mathbb{R}^{n+1} .

The second fundamental B of the immersion is defined by

$$B(Y, Z) = \langle \bar{\nabla}_Y \nu, Z \rangle$$

and the mean curvature H by

$$H = \frac{1}{n} \text{tr}(B).$$

Now we recall the following well-known identity

$$(3) \quad \frac{1}{2} \Delta |X|^2 = nH \langle \nu, X \rangle - n,$$

where X is the position vector.

We finish by recalling the definition of higher order mean curvatures. They are extrinsic geometric invariants defined from the second fundamental form and generalizing the mean curvature. We saw that

$$H = \frac{1}{n} \sigma_1(\kappa_1, \dots, \kappa_n),$$

where σ_1 is the first symmetric polynomial and $\kappa_1, \dots, \kappa_n$ the principal curvatures of M . The higher order mean curvatures are defined for $k \in \{1, \dots, n\}$ by

$$H_k = \frac{1}{\binom{n}{k}} \sigma_k(\kappa_1, \dots, \kappa_n),$$

where σ_k is the k -th symmetric polynomial, that is,

$$\sigma_k(x_1, \dots, x_n) = \sum_{1 \leq i_1, \dots, i_k \leq n} x_{i_1} \cdots x_{i_k}.$$

This definition is equivalent to

$$H_k = \frac{1}{k! \binom{n}{k}} \sum_{\substack{1 \leq i_1, \dots, i_k \leq n \\ 1 \leq j_1, \dots, j_k \leq n}} \epsilon \left(\begin{matrix} i_1, \dots, i_k \\ j_1, \dots, j_k \end{matrix} \right) B_{i_1 j_1} \cdots B_{i_k j_k},$$

where the B_{ij} 's are the coefficients of the second fundamental form B in a local orthonormal frame $\{e_1, \dots, e_n\}$. Moreover, we denote by $\epsilon \left(\begin{matrix} i_1, \dots, i_k \\ j_1, \dots, j_k \end{matrix} \right)$ the usual symbols for permutation. Finally, by convention, we set $H_0 = 1$

and $H_{n+1} = 0$.

These mean curvatures satisfy some properties as the Hsiung-Minkowski formula (see [5])

$$(4) \quad \int_M (H_{k-1} - H_k \langle X, \nu \rangle) = 0,$$

and the following inequalities

Lemma 1.1. *If $k \in \{1, \dots, n\}$, and H_k is a positive function, then*

$$H_k^{\frac{1}{k}} \leq H_{k-1}^{\frac{1}{k-1}} \leq \dots \leq H_2^{\frac{1}{2}} \leq H.$$

2 An L^2 -approach to the problem

We prove Theorems 1 and 2 in two steps. First, we prove that if the pinching condition (P_C) is satisfied, then M is close to a sphere in an L^2 -sense. For this, we prove a first lemma which gives an estimate of the L^2 -norm of the position vector X .

Lemma 2.1. *If the pinching condition (P_C) is satisfied for $C < \frac{n}{2} \|H_k\|_{2p}^2$, then*

$$\frac{n\lambda_1(M) \left(\int_M H_{k-1}\right)^4}{\left(C + \lambda_1(M) \left(\int_M H_{k-1}\right)^2\right)^2} \leq \|X\|_2^2 \leq \frac{n}{\lambda_1(M)} \leq A_1,$$

where A_1 is a positive constant depending only on n , $\|H\|_\infty$ and $\|H_k\|_{2p}$.

Proof: If (P_C) is satisfied, we have:

$$\lambda_1(M) \left(\int_M H_{k-1}\right)^2 \geq n\|H_k\|_{2p}^2 - C.$$

If, in addition, we assume that $C < \frac{n}{2} \|H_k\|_{2p}^2$, we get

$$\lambda_1(M) \left(\int_M H_{k-1}\right)^2 \geq \frac{n}{2} \|H_k\|_{2p}^2,$$

and so

$$(5) \quad \frac{n}{\lambda_1(M)} \leq \frac{2 \left(\int_M H_{k-1}\right)^2}{\|H_k\|_{2p}^2} \leq \frac{2\|H\|_\infty^{2(k-1)}}{\|H_k\|_{2p}^2}.$$

Moreover, by the variational characterization of $\lambda_1(M)$, we have

$$\lambda_1(M) \int_M |X|^2 \leq \int_M \left(\sum_{i=1}^{n+1} |dX_i|^2 \right) = n,$$

where X_i are the functions defined by $X = \sum_{i=1}^{n+1} X_i \partial_i$, where $\{\partial_1, \dots, \partial_{n+1}\}$ is the canonical frame of \mathbb{R}^{n+1} . So we have $\|X\|_2^2 \leq \frac{n}{\lambda_1(M)}$, and by (5),

$$\|X\|_2^2 \leq A_1(n, \|H\|_\infty, \|H_k\|_{2p}).$$

For the left hand side, we have

$$\begin{aligned} \lambda_1(M) \left(\int_M |X|^2 \right) \left(\int_M H_{k-1} \right)^4 &\leq n \left(\int_M H_{k-1} \right)^4 \\ &\leq n \left(\int_M H_k \langle X, \nu \rangle \right)^4 \\ &\leq n \left(\int_M H_k^2 \right)^2 \left(\int_M |X|^2 \right)^2. \end{aligned}$$

Then, by the Hölder inequality, we deduce

$$\lambda_1(M) \left(\int_M H_{k-1} \right)^4 \leq n \|H_k\|_{2p}^2 \left(\int_M |X|^2 \right),$$

and with the pinching condition,

$$\|X\|_2^2 \geq \frac{n \lambda_1(M) \left(\int_M H_{k-1} \right)^4}{\left(C + \lambda_1(M) \left(\int_M H_{k-1} \right)^2 \right)^2}.$$

□

From now, we denote by X^T the orthogonal projection of X on M . That is, if for $x \in M$, $\{e_1, \dots, e_n\}$ is an orthonormal frame of $T_x M$, then

$$X^T = \sum_{i=1}^n \langle X, e_i \rangle e_i = X - \langle X, \nu \rangle \nu.$$

In the following lemma, we show that the pinching condition (P_C) implies that the L^2 -norm of X^T is close to zero.

Lemma 2.2. *The pinching condition (P_C) with $C < \frac{n}{2} \|H_k\|_{2p}^2$ implies*

$$\|X^T\|_2^2 \leq A_2 C,$$

where A_2 is a positive constant depending only on n , $\|H\|_\infty$ and $\|H_k\|_{2p}$.

Proof: We saw that

$$\lambda_1(M) \int_M |X|^2 \leq n,$$

so by the Hsiung-Minkowski formula and the Cauchy-Schwarz inequality

$$\begin{aligned} \lambda_1(M) \int_M |X|^2 \left(\int_M H_{k-1} \right)^2 &\leq n \left(\int_M H_{k-1} \right)^2 \\ &\leq n \left(\int_M H_k \langle X, \nu \rangle \right)^2 \\ &\leq \|H_k\|_2^2 \int_M \langle X, \nu \rangle^2 \\ &\leq \|H_k\|_{2p}^2 \int_M \langle X, \nu \rangle^2. \end{aligned}$$

Then we deduce

$$\begin{aligned} n \|H_k\|_{2p}^2 \|X^T\|_2^2 &\leq n \|H_k\|_{2p}^2 \left(\int_M (|X|^2 - \langle X, \nu \rangle^2) \right) \\ &\leq n \|H_k\|_{2p}^2 \left[\int_M |X|^2 - \lambda_1(M) \left(\int_M H_{k-1} \right)^2 \int_M |X|^2 \right] \\ &\leq \left[n \|H_k\|_{2p}^2 - \lambda_1(M) \left(\int_M H_{k-1} \right)^2 \right] \|X\|_2^2 \\ &\leq C \|X\|_2^2 \leq A_1 C. \end{aligned}$$

Finally, we get

$$\|X^T\|_2^2 \leq \frac{A_1 C}{n \|H_k\|_{2p}^2} = A_2 C.$$

□

In order to prove assertion *i*) of Theorem 1, we will show that

$$\left\| |X| - \sqrt{\frac{n}{\lambda_1(M)}} \right\|_\infty \leq \varepsilon.$$

For this, we need an upper bound on the L^2 -norm of the function

$$\varphi := |X| \left(|X| - \sqrt{\frac{n}{\lambda_1(M)}} \right)^2.$$

Before getting such an estimate, we introduce the two following vector fields:

$$\begin{cases} Y = nH_k\nu - \lambda_1(M) \left(\int_M H_{k-1} \right) X, \\ Z = \sqrt{\frac{n}{\lambda_1}} \frac{|X|^{1/2} H_k}{\left(\int_M H_{k-1} \right)} \nu - \frac{X}{|X|^{1/2}}. \end{cases}$$

First, we have the following:

Lemma 2.3. *The pinching condition (P_C) implies*

$$\|Y\|_2^2 \leq nC.$$

Proof: We have

$$\begin{aligned} \|Y\|_2^2 &= n^2 \int_M H_k^2 + \lambda_1(M) \left(\int_M H_{k-1} \right)^2 \int_M |X|^2 \\ &\quad - 2n\lambda_1(M) \left(\int_M H_{k-1} \right) \int_M H_k \langle X, \nu \rangle \\ &\leq n^2 \|H_k\|_{2p}^2 + n\lambda_1(M) \left(\int_M H_{k-1} \right)^2 - 2n\lambda_1(M) \left(\int_M H_{k-1} \right)^2 \\ &\leq n \left(n \|H_k\|_{2p}^2 - n\lambda_1(M) \left(\int_M H_{k-1} \right)^2 \right) \\ &\leq nC, \end{aligned}$$

where we used the Hsiung-Minkowski formula (4), and the fact that

$$\|X\|_2^2 \leq \frac{n}{\lambda_1(M)}.$$

□

We also have

Lemma 2.4. *If the pinching condition (P_C) is satisfied, with $C < \frac{n}{2} \|H_k\|_{2p}^2$, then*

$$\|Z\|_2^2 \leq A_3 C,$$

where A_3 is a positive constant depending only on n , $\|H\|_\infty$ and $\|H_k\|_{2p}$.

Proof: We have

$$\begin{aligned} \|Z\|_2^2 &= \frac{n}{\lambda_1(M) \left(\int_M H_{k-1} \right)^2} \int_M |X| H_k^2 + \int_M |X| - 2 \frac{\sqrt{\frac{n}{\lambda_1(M)}}}{\int_M H_{k-1}} \int_M H_k \langle X, \nu \rangle \\ &\leq \frac{n}{\lambda_1(M) \left(\int_M H_{k-1} \right)^2} \int_M |X| H_k^2 + \int_M |X| - 2 \sqrt{\frac{n}{\lambda_1(M)}} \end{aligned}$$

By the Hölder inequality, we get

$$\begin{aligned}
\|Z\|_2^2 &\leq \frac{n}{\lambda_1(M) \left(\int_M H_{k-1}\right)^2} \left(\int_M H_k^4\right)^{1/2} \left(\int_M |X|^2\right)^{1/2} \\
&\quad + \left(\int_M |X|^2\right)^{1/2} - 2\sqrt{\frac{n}{\lambda_1(M)}} \\
&\leq \sqrt{\frac{n}{\lambda_1(M)}} \left[\frac{n}{\lambda_1(M) \left(\int_M H_{k-1}\right)^2} \|H_k\|_{2p}^2 - 1 \right] \\
&\leq \left(\frac{n}{\lambda_1(M)}\right)^{3/2} \frac{1}{\left(\int_M H_{k-1}\right)^2} \left[n\|H_k\|_{2p}^2 - \lambda_1(M) \left(\int_M H_{k-1}\right)^2 \right] \\
&\leq A_3 C,
\end{aligned}$$

where A_3 depends only on n , $\|H\|_\infty$ and $\|H_k\|_{2p}$. Note that we have used Lemma 1.1 and the fact that $\frac{n}{\lambda_1(M)} \leq \frac{2\|H\|_\infty^{2(k-1)}}{\|H_k\|_{2p}^2}$. \square

Now, we give an upper bound for the L^2 -norm of the function φ .

Lemma 2.5. *The pinching condition (P_C) with $C < \frac{n}{2}\|H_k\|_{2p}^2$ implies*

$$\|\varphi\|_2 \leq A_4 \|\varphi\|_\infty^{3/4} C^{1/4}.$$

Proof: We have

$$\|\varphi\|_2 = \left(\int_M \varphi^{3/2} \varphi^{1/2}\right)^{1/2} \leq \|\varphi\|_\infty^{3/4} \|\varphi^{1/2}\|_1^{1/2}.$$

Moreover,

$$\begin{aligned}
\int_M \varphi^{1/2} &= \left\| |X|^{1/2} X - \sqrt{\frac{n}{\lambda_1(M)}} \frac{X}{|X|^{1/2}} \right\|_1 \\
&= \left\| -\frac{|X|^{1/2}}{\lambda_1(M) \int_M H_{k-1}} Y + \frac{n}{\lambda_1(M) \int_M H_{k-1}} \nu - \sqrt{\frac{n}{\lambda_1(M)}} \frac{X}{|X|^{1/2}} \right\|_1 \\
&\leq \left\| \frac{|X|^{1/2}}{\lambda_1(M) \int_M H_{k-1}} Y \right\|_1 + \sqrt{\frac{n}{\lambda_1(M)}} \|Z\|_1.
\end{aligned}$$

By the Hölder inequality, we get

$$\begin{aligned}
\left\| \frac{|X|^{1/2}}{\lambda_1(M) \int_M H_{k-1}} Y \right\|_1 &\leq \frac{1}{\lambda_1} \left(\int_M |X|^2\right)^{1/4} \|Y\|_2 \\
&\leq \frac{A_1^{3/4}}{n^{1/2}} C^{1/2}.
\end{aligned}$$

Finally, from Lemmas 2.3 and 2.4, we obtain

$$\|\varphi^{1/2}\|_1^{1/2} \leq A_4 C^{1/4},$$

where A_4 is a positive constant depending only on n , $\|H\|_\infty$ and $\|H_k\|_{2p}$.
 \square

3 Proof of Theorem 1

The proof of Theorem 1 is an immediate consequence of the two following lemmas:

Lemma 3.1. *For $p \geq 2$ and any $\eta > 0$, there exists $K_\eta(n, \|H\|_\infty, \|H_k\|_{2p})$ so that if (P_{K_η}) is true, then $\|\varphi\|_\infty \leq \eta$. Moreover, $K_\eta \rightarrow 0$ when $\|H\|_\infty \rightarrow \infty$ or $\eta \rightarrow 0$.*

Lemma 3.2. *[Colbois-Grosjean [2]] Let x_0 be a point of the sphere $S(0, R)$ in \mathbb{R}^{n+1} with the center at the origin and of radius R . Assume that $x_0 = Re$ with $e \in \mathbb{S}^n$. Now let (M^n, g) be a compact, connected, oriented n -dimensional Riemannian manifold without boundary isometrically immersed in \mathbb{R}^{n+1} . If the image of M is contained in $(B(0, R + \eta) \setminus B(0, R - \eta)) \setminus B(x_0, \rho)$ with $\rho = 4(2n - 1)\eta$. Then there exists a point $y_0 \in M$ so that the mean curvature of M in y_0 satisfies $|H(y_0)| \geq \frac{1}{4n\eta}$.*

We will prove Lemma 3.1 in Section 6. Now, we will prove Theorem 1 by using these two lemmas.

Proof of Theorem 1 Let $\varepsilon > 0$ and consider the function

$$f(t) := t \left(t - \sqrt{\frac{n}{\lambda_1(M)}} \right)^2.$$

We set

$$\eta(\varepsilon) := \inf \left\{ f \left(\sqrt{\frac{n}{\lambda_1(M)}} - \varepsilon \right), f \left(\sqrt{\frac{n}{\lambda_1(M)}} + \varepsilon \right), \frac{1}{27\|H\|_\infty^3} \right\}.$$

By definition, $\eta(\varepsilon) > 0$, and by Lemma 3.1, there exists $K_{\eta(\varepsilon)}$ such that for all $x \in M$,

$$(6) \quad f(|X|(x)) \leq \eta(\varepsilon).$$

Now to prove the theorem, it is sufficient to assume $\varepsilon < \frac{2}{3\|H\|_\infty}$. We will show that either

$$(7) \quad \sqrt{\frac{n}{\lambda_1(M)}} - \varepsilon \leq |X| \leq \sqrt{\frac{n}{\lambda_1(M)}} + \varepsilon \quad \text{or} \quad |X| < \frac{1}{3}\sqrt{\frac{n}{\lambda_1(M)}}$$

By examining the function f , it is easy to see that f has a unique local maximum at $\frac{1}{3}\sqrt{\frac{n}{\lambda_1(M)}}$. Moreover, from the definition of $\eta(\varepsilon)$, we have

$$\eta(\varepsilon) < \frac{4}{27\|H\|_\infty^3} \leq \frac{4}{27} \left(\frac{n}{\lambda_1(M)} \right)^{3/2} = f \left(\frac{1}{3}\sqrt{\frac{n}{\lambda_1(M)}} \right).$$

Since we assume $\varepsilon < \frac{2}{3\|H\|_\infty} \leq \frac{2}{3}\sqrt{\frac{n}{\lambda_1(M)}}$, we have

$$\frac{1}{3}\sqrt{\frac{n}{\lambda_1(M)}} < \sqrt{\frac{n}{\lambda_1(M)}} - \varepsilon,$$

which with (6) yields (7).

Now, from Lemma 2.1, we deduce that there exists a point $y_0 \in M$ such that

$$|X(y_0)|^2 \geq \frac{n\lambda_1(M) \left(\int_M H_{k-1} \right)^4}{\left(K_{\eta(\varepsilon)} + \lambda_1(M) \left(\int_M H_{k-1} \right)^2 \right)^2}.$$

Since $K_{\eta(\varepsilon)} < \frac{n}{2}\|H_k\|_{2p}^2$, the condition (P_C) implies

$$K_{\eta(\varepsilon)} < \frac{n}{2}\|H_k\|_{2p}^2 \leq \lambda_1(M) \left(\int_M H_{k-1} \right)^2 \leq 2\lambda_1(M) \left(\int_M H_{k-1} \right)^2.$$

We deduce that

$$|X(y_0)| \geq \frac{1}{3}\sqrt{\frac{n}{\lambda_1(M)}}.$$

Since M is connected, for any $x \in M$,

$$\sqrt{\frac{n}{\lambda_1(M)}} - \varepsilon \leq |X|(x) \leq \sqrt{\frac{n}{\lambda_1(M)}} + \varepsilon,$$

which proves the assertion *i*) of the theorem.

In order to prove the second, we consider the pinching condition (P_{C_ε}) with $C_\varepsilon = K_{\eta(\frac{\varepsilon}{4(2n-1)})}$. Then assertion *i*) is still valid.

Let $x = \sqrt{\frac{n}{\lambda_1(M)}} e \in S \left(0, \sqrt{\frac{n}{\lambda_1(M)}} \right)$, with $e \in \mathbb{S}^n$ and assume that

$B(x, \varepsilon) \cap M = \emptyset$. We can apply Lemma 3.2. So, there exists a point $y_0 \in M$ such that $|H(y_0)| \geq \frac{2n-1}{n\varepsilon} > \|H\|_\infty$ since we assumed $\varepsilon < \frac{2}{3\|H\|_\infty} \leq \frac{2n-1}{n\|H\|_\infty}$. This is a contradiction and so $B(x, \varepsilon) \cap M \neq \emptyset$. The assertion *ii*) is satisfied and $C_\varepsilon \rightarrow 0$ when $\|H\|_\infty \rightarrow \infty$ or $\varepsilon \rightarrow 0$. \square

4 Proof of Theorem 2

From Theorem 1, we know that for any $\varepsilon > 0$, there exists C_ε depending only on n , $\|H\|_\infty$ and $\|H_k\|_{2p}$ so that if (P_{C_ε}) is true, then

$$\left| |X|(x) - \sqrt{\frac{n}{\lambda_1(M)}} \right| \leq \varepsilon$$

for all $x \in M$. Since $\sqrt{n}\|H\|_\infty \leq \|B\|_\infty$, it is easy to see that we can assume that C_ε depends only on n , $\|B\|_\infty$ and $\|H_k\|_{2p}$.

The proof of Theorem 2 is an immediate consequence of the following lemma about the L^∞ -norm of X^T .

Lemma 4.1. *For $p \geq 2$ and any $\eta > 0$, there exists $K_\eta(n, \|B\|_\infty, \|H_k\|_{2p})$ so that if (P_{K_η}) is true, then $\|X^T\|_\infty \leq \eta$.*

We will prove this lemma in Section 6.

Proof of Theorem 2 Let $\varepsilon < \frac{1}{2}\sqrt{\frac{n}{\|B\|_\infty}} \leq \sqrt{\frac{n}{\lambda_1(M)}}$. This choice of ε implies that if the pinching condition (P_{C_ε}) is true, then $|X|$ never vanishes, and so we can consider the following map

$$\begin{aligned} F : M &\longrightarrow S\left(0, \sqrt{\frac{n}{\lambda_1(M)}}\right) \\ x &\longmapsto \sqrt{\frac{n}{\lambda_1(M)}} \frac{X}{|X|}. \end{aligned}$$

Without any pinching condition, a straightforward computation yields to

$$(8) \quad \left| |dF_x(u)|^2 - 1 \right| \leq \left| \frac{n}{\lambda_1(M)} \frac{1}{|X|^2} - 1 \right| + \frac{n}{\lambda_1(M)} \frac{1}{|X|^4} \langle u, X \rangle^2,$$

for any unitary vector $u \in T_x M$. But,

$$\left| \frac{n}{\lambda_1(M)} \frac{1}{|X|^2} - 1 \right| = \frac{1}{|X|^2} \left| \frac{n}{\lambda_1(M)} - |X|^2 \right| \leq \varepsilon \frac{\sqrt{\frac{n}{\lambda_1(M)}} + |X|}{|X|^2} \leq \varepsilon \frac{2\sqrt{\frac{n}{\lambda_1(M)}} + \varepsilon}{\left(\sqrt{\frac{n}{\lambda_1(M)}} - \varepsilon\right)^2}$$

We recall that $\frac{n}{A_1} \leq \lambda_1 \leq \|B\|_\infty^2$. Since we assume $\varepsilon < \frac{1}{2} \sqrt{\frac{n}{\|B\|_\infty}}$, the right hand side is bounded by a constant depending only on n , $\|B\|_\infty$ and $\|H_k\|_{2p}$. So we have

$$(9) \quad \left| \frac{n}{\lambda_1(M)} \frac{1}{|X|^2} - 1 \right| \leq \varepsilon \gamma(n, \|B\|_\infty, \|H_k\|_{2p}).$$

Moreover, since $C_\varepsilon \rightarrow 0$ when $\varepsilon \rightarrow 0$, there exists $\varepsilon(n, \|B\|_\infty, \|H_k\|_{2p}, \eta)$ so that $C_\varepsilon \leq K_\eta$ (where K_η is the constant of Lemma 4.1) and so, $\|X^T\|_\infty \leq \eta$. As before, there exists a constant δ depending on n , $\|B\|_\infty$ and $\|H_k\|_{2p}$ such that

$$(10) \quad \frac{n}{\lambda_1(M)} \frac{1}{|X|^4} \langle u, X \rangle^2 \leq \frac{n}{\lambda_1(M)} \frac{1}{|X|^4} \|X^T\|_\infty^2 \leq \eta^2 \delta(n, \|B\|_\infty, \|H_k\|_{2p}).$$

Then, from (8), (9) and (10), we deduce that (P_{C_ε}) implies

$$\left| |dF_x(u)|^2 - 1 \right| \leq \varepsilon \gamma + \eta^2 \delta.$$

Take $\eta = \sqrt{\frac{\theta}{2\delta}}$. We can assume that ε is small enough to have $\varepsilon \gamma \leq \frac{\theta}{2}$. In that case, we have

$$\left| |dF_x(u)|^2 - 1 \right| \leq \theta.$$

Now, it is sufficient to fix $\theta \in]0, 1[$ and, F is a local diffeomorphism from M into $S\left(0, \sqrt{\frac{n}{\lambda_1(M)}}\right)$. Since $S\left(0, \sqrt{\frac{n}{\lambda_1(M)}}\right)$ is simply connected for $n \geq 2$, the map F is a global diffeomorphism. \square

5 Application to almost-Einstein Hypersurfaces

In this section, we give an application of Theorems 1 and 2 to almost Einstein hypersurfaces. In fact, we obtain two different rigidity results.

Corollary 1. *Let (M^n, g) be a connected, oriented Riemannian manifold without boundary isometrically immersed in \mathbb{R}^{n+1} . If (M^n, g) is almost-Einstein, that is, $\|\text{Ric} - kg\|_\infty \leq \varepsilon$ for a positive constant k , with ε small enough depending on n , k and $\|H\|_\infty$, then*

$$d_H \left(M, \mathbb{S}^n \left(\sqrt{\frac{n-1}{k}} \right) \right) \leq \varepsilon.$$

Corollary 2. *Let (M^n, g) be a compact, connected, oriented Riemannian manifold without boundary isometrically immersed in \mathbb{R}^{n+1} . If (M^n, g) is almost-Einstein, that is, $\|\text{Ric} - kg\|_\infty \leq \varepsilon$ for a positive constant k , with ε small enough depending on n , k and $\|B\|_\infty$, then M is diffeomorphic and almost isometric to $\mathbb{S}^n \left(\sqrt{\frac{n-1}{k}} \right)$*

Proof: Assume that $k = n - 1$. By the assumption $\|\text{Ric} - (n - 1)g\|_\infty \leq \varepsilon$, the Lichnerowicz theorem implies that

$$\lambda_1(M) \geq \frac{n(n - 1 - \varepsilon)}{n - 1} = n - \frac{n\varepsilon}{n - 1}.$$

So, for $p \geq 2$, we have

$$\begin{aligned} \lambda_1(M) \left(\int_M H \right)^2 - n \|H_2\|_{2p}^2 &\geq n \left(1 - \frac{\varepsilon}{n - 1} \right) \left(\int_M H_2^{1/2} \right)^2 - n \|H_2\|_{2p}^2 \\ &\geq n \left(1 - \frac{\varepsilon}{n - 1} \right) \inf \{H_2\} - n \text{Sup} \{H_2\} \\ &\geq n \left(1 - \frac{\varepsilon}{n - 1} \right)^2 - n \left(1 + \frac{\varepsilon}{n - 1} \right) \\ &\geq \frac{n\varepsilon^2}{(n - 1)^2} - \frac{3n\varepsilon}{n - 1} = -\beta_n(\varepsilon), \end{aligned}$$

where β_n is a positive function such that $\beta_n(\varepsilon) \rightarrow 0$ when $\varepsilon \rightarrow 0$.

We can choose ε small enough to have $\beta_n(\varepsilon) \leq C(n, \|B\|_\infty, \|H_2\|_{2p})$ of Theorem 2, and we deduce that there exists ε depending only on n and $\|B\|_\infty$ so that if $\|\text{Ric} - (n - 1)g\|_\infty \leq \varepsilon$, then M is diffeomorphic and almost isometric to \mathbb{S}^n . Since $1 - \frac{n}{\varepsilon} \leq H_2 \leq 1 + \frac{n}{\varepsilon}$, there is no dependence on $\|H_2\|_{2p}$. By homothety, we get the result for any $k > 0$.

The proof of Corollary 1 is the same, we use Theorem 1 instead of Theorem 2. \square

As we mentioned, these two corollaries are to be compared to the following theorem obtained by a pinching result associated with the Lichnerowicz inequality.

Theorem 3. *Let (M^n, g) be a compact, connected, oriented Riemannian manifold without boundary isometrically immersed in \mathbb{R}^{n+1} and $p > \frac{n}{2}$. Then for any $k > 0$, there exists $\varepsilon(k, n, \|K\|_{2p})$ (where K is the sectional curvature of M) such that if*

$$\|\text{Ric} - kg\|_\infty \leq \varepsilon,$$

then M is homeomorphic to \mathbb{S}^n .

Proof: The assumption

$$\|\text{Ric} - kg\|_\infty \leq \varepsilon,$$

implies that the scalar curvature satisfies

$$0 < n(k - \varepsilon) \leq \text{Scal} \leq n(k + \varepsilon).$$

So the first eigenvalue of the Laplacian can be bounded from above

$$\lambda_1(M) \leq \frac{n(k + \varepsilon)}{n - 1}.$$

On the other hand, the Lichnerowicz theorem says that

$$\lambda_1(M) \geq \frac{n(k - \varepsilon)}{n - 1}.$$

Now, let us recall the following theorem due to E. Aubry ([1]), which a generalization of a theorem of Ilias ([6]).

Theorem (Aubry [1]). *Let p, R and A be some real numbers such that $p > \frac{n}{2}$, $R > 0$ and $A > 0$. Let (M^n, g) be a complete Riemannian manifold. There exists $\alpha(p, n, A) > 0$ such that if*

$$\text{Sup}_x \frac{\|(\underline{\text{Ric}} - (n - 1))^- \|_{L^p(B(x, R))}}{V(B(x, R))} \leq \alpha(p, n, A),$$

$\|K\|_{2p} \leq A$, and

$$\lambda_1(M) \leq n(1 + \alpha(p, n, A)),$$

then M is homeomorphic to \mathbb{S}^n .

In this theorem, $\underline{\text{Ric}}(x)$ is the smallest eigenvalue of the symmetric bilinear form $\text{Ric}(x)$ on $T_x M$, and $(\underline{\text{Ric}} - (n - 1))^- = \max(0, -\underline{\text{Ric}} + (n - 1))$.

Since M is almost-Einstein, we are precisely in the assumptions of this theorem, and it is sufficient to choose $\varepsilon(k, n, \|K\|_{2p}) > 0$ small enough. \square

6 Proof of the technical Lemmas

The proof of Lemmas 3.1 and 4.1 is based on the following result due to Colbois and Grosjean [2] using a Nirenberg-Moser type of argument.

Lemma 6.1. *Let (M^n, g) be a compact, connected, oriented Riemannian manifold without boundary isometrically immersed in \mathbb{R}^{n+1} and let ξ be a nonnegative continuous function on M such that ξ^k is smooth for $k \geq 2$. Assume there exist $0 \leq l < m \leq 2$ such that*

$$\frac{1}{2}\xi^{2k-2}\Delta\xi^2 \leq \operatorname{div} \omega + (\alpha_1 + k\alpha_2)\xi^{2k-l} + (\beta_1 + k\beta_2)\xi^{2k-m},$$

where ω is a 1-form and $\alpha_1, \alpha_2, \beta_1$ and β_2 are some nonnegative constants. Then for any $\eta > 0$, there exists a positive constant L depending on $n, \delta, \alpha_1, \alpha_2, \beta_1, \beta_2, \|H\|_\infty$ and η so that if $\|\xi\|_\infty > \eta$, then

$$\|\eta\|_\infty \leq L\|\xi\|_2.$$

Moreover, L is bounded when $\eta \rightarrow +\infty$ and if $\beta_1 > 0$, then $L \rightarrow +\infty$ if $\|H\|_\infty \rightarrow +\infty$ or $\eta \rightarrow 0$.

In order to prove 3.1 and 4.1, it is sufficient to find an upper bound for the functions

$$\begin{cases} \varphi^{2k-2}\Delta\varphi^2 \\ |X^T|^{2k-2}\Delta|X^T|^2. \end{cases}$$

For this, the pinching condition (P_C) is used only one time, to obtain an upper bound of $\|X\|_\infty$ depending only on $n, \|H\|_\infty$ and $\|H_k\|_{2p}$.

Lemma 6.2. *If the pinching condition (P_C) is satisfied with $C < \frac{n}{2}\|H_k\|_{2p}^2$, then there exists $E(n, \|H\|_\infty, \|H_k\|_{2p})$ such that $\|X\|_\infty \leq E$.*

Proof: From (3), we have

$$\frac{1}{2}\Delta|X|^2|X|^{2l-2} \leq n\|H\|_\infty|X|^{2l-1}.$$

We apply Proposition 6.1 to the function $\xi = |X|$. We now that if $\|X\|_\infty > E$, then there exists a constant $L(n, \|H\|_\infty, E)$ such that

$$\|X\|_\infty \leq L\|X\|_2.$$

By the pinching condition (P_C) with $C < \frac{n}{2}\|H_k\|_{2p}^2$, we obtain from Lemma 2.1 that

$$\|X\|_\infty \leq LA_1(n, \|H\|_\infty, \|H_k\|_{2p})^{1/2}.$$

But, L is bounded when $E \rightarrow 0$, so we can choose $E = E(n, \|H\|_\infty, \|H_k\|_{2p})$ big enough to have

$$LA_1(n, \|H\|_\infty, \|H_k\|_{2p})^{1/2} < E.$$

In that case, we have $\|X\|_\infty \leq E(n, \|H\|_\infty, \|H_k\|_{2p})$. \square

Then, the proof of Lemmas 3.1 and 4.1 is exactly the same as the proof of the technical Lemmas in [2], [8] or [9].

References

- [1] E. Aubry, *Variétés de courbure de Ricci presque minorée: inégalités géométriques optimales et stabilité des variétés extrémales*, Ph.D. thesis, Université Joseph Fourier, Grenoble, 2003.
- [2] B. Colbois and J.F. Grosjean, *A pinching theorem for the first eigenvalue of the Laplacian on hypersurfaces of the Euclidean space*, Comment. Math. Helv. **82** (2007), 175–195.
- [3] A. Fialkow, *Hypersurfaces of a space of constant curvature*, Ann. of Math. **39** (1938), 762–783.
- [4] J.F. Grosjean, *Upper bounds for the first eigenvalue of the Laplacian on compact manifolds*, Pac. J. Math. **206** (2002), no. 1, 93–111.
- [5] C.C. Hsiung, *Some integral formulae for closed hypersurfaces*, Math. Scand. **2** (1954), 286–294.
- [6] S. Ilias, *Un nouveau résultat de pincement de la première valeur propre du Laplacien et conjecture du diamètre pincé*, Ann. Inst. Fourier **43** (1993), no. 3, 843–863.
- [7] R.C. Reilly, *On the first eigenvalue of the Laplacian for compact submanifolds of Euclidean space*, Comment. Math. Helv. **52** (1977), 525–533.
- [8] J. Roth, *Extrinsic radius pinching for hypersurfaces of space forms*, To appear in Diff. Geom. Appl.
- [9] ———, *Extrinsic radius pinching in space forms of nonnegative sectional curvature*, To appear in Math. Z.
- [10] T.Y. Thomas, *On closed space of constant mean curvature*, Amer. J. Math. **58** (1936), 702–704.